arXiv:0806.4708v4 [math.DG] 11 Feb 2009

KÄHLER MANIFOLDS WITH QUASI-CONSTANT HOLOMORPHIC CURVATURE.

Włodzimierz Jelonek

Abstract. The aim of this paper is to classify compact Kähler manifolds with quasi-constant holomorphic sectional curvature.

0. Introduction. The aim of the present paper is to classify compact, simply connected Kähler manifolds (M, g, J) admitting a global, 2-dimensional, J-invariant distribution $\mathcal{D}$ having the following property: The holomorphic curvature $K(\pi) = R(X, JX, JX, X)$ of any J-invariant 2-plane $\pi \subset T_xM$, where $X \in \pi$ and $g(X, X) = 1$, depends only on the point x and the number $|X_{\mathcal{D}}| = \sqrt{g(X_{\mathcal{D}}, X_{\mathcal{D}})}$, where $X_{\mathcal{D}}$ is an orthogonal projection of X on $\mathcal{D}$. In this case we have

$$R(X, JX, JX, X) = \phi(x, |X_{\mathcal{D}}|)$$

where $\phi(x, t) = a(x) + b(x)t^2 + c(x)t^4$ and a, b, c are smooth functions on M. Also $R = a\Pi + b\Phi + c\Psi$ for certain curvature tensors $\Pi, \Phi, \Psi \in \bigotimes^4 \mathfrak{X}^*(M)$ of Kähler type. The investigation of such manifolds, called QCH Kähler manifolds, was started by G. Ganchev and V. Mihova in [G-M-1],[G-M-2]. In our paper we shall use their local results to obtain a global classification of such manifolds under the assumption that $\dim M = 2n \geq 6$. By $\mathcal{E}$ we shall denote the $(\dim M - 2)$-dimensional distribution which is the orthogonal complement of $\mathcal{D}$ in TM. If X is a local unit section of $\mathcal{D}$ then $\{X, JX\}$ is a local orthonormal basis of $\mathcal{D}$ and the function $\kappa = \sqrt{(div_{\mathcal{E}} X)^2 + (div_{\mathcal{E}} JX)^2}$ does not depend on the choice of X. We classify compact, simply connected QCH Kähler manifolds satisfying the conditions int $B = \emptyset$ and $U \neq \emptyset$ where $B = \{x \in U : b(x) = 0\}, U = \{x \in M : \kappa(x) \neq 0\}$. First we show that (M, g, J) admits a global holomorphic Killing vector field with a Killing potential, which is a special Kähler-Ricci potential. Next we use the results of Derdziński and Maschler [D-M-1], who classified compact Kähler manifolds admitting special Kähler-Ricci potentials. As a corollary we prove that every compact, simply connected QCH Kähler manifold with $\kappa \neq 0$ and an analytic Riemannian metric g is a holomorphic $\mathbb{CP}^1$-bundle over $\mathbb{CP}^{n-1}$ (with $\mathcal{D}$ being an integrable distribution whose leaves are the $\mathbb{CP}^1$ fibers of the bundle) or it is $\mathbb{CP}^n$ with a metric of constant holomorphic sectional curvature (in this case $\mathcal{D}$ is any J-invariant 2-dimensional distribution on $\mathbb{CP}^n$ with $\kappa \neq 0$; however, such distributions may not exist).

MS Classification: 53C55,53C25. Key words and phrases: Kähler manifold, holomorphic sectional curvature, quasi constant holomorphic sectional curvature, special Kähler-Ricci potential

Typeset by $\mathcal{A}\mathcal{M}\mathcal{S}$-TEX

1. Special frame. Let (M,g,J) be a $2n$-dimensional Kähler manifold with a 2-dimensional J-invariant distribution $\mathcal{D}$. Let $\mathfrak{X}(M)$ denote the algebra of all differentiable vector fields on M and $\Gamma(\mathcal{D})$ denote the set of local sections of the distribution $\mathcal{D}$. If $X \in \mathfrak{X}(M)$ then by $X^\flat$ we shall denote the 1-form $\phi \in \mathfrak{X}^*(M)$ dual to X with respect to g, i.e. $\phi(Y) = X^\flat(Y) = g(X,Y)$. By Ω we shall denote the Kähler form of (M,g,J) i.e. $\Omega(X,Y) = g(JX,Y)$. Let us denote by $\mathcal{E}$ the distribution $\mathcal{D}^\perp$, which is a $2(n-1)$-dimensional, J-invariant distribution. By h, m respectively we shall denote the tensors $h = g \circ (p_{\mathcal{D}} \times p_{\mathcal{D}}), m = g \circ (p_{\mathcal{E}} \times p_{\mathcal{E}})$, where $p_{\mathcal{D}}, p_{\mathcal{E}}$ are the orthogonal projections on $\mathcal{D}, \mathcal{E}$ respectively. It follows that $g = h + m$. By ω we shall denote the Kähler form of $\mathcal{D}$ i.e. $\omega(X,Y) = h(JX,Y)$ and by Ω_m the Kähler form of $\mathcal{E}$ i.e. $\Omega_m(X,Y) = m(JX,Y)$. For any local section $X \in \Gamma(\mathcal{D})$ we define $div_{\mathcal{E}} X = tr_m \nabla X^\flat = m^{ij} \nabla_{e_i} X^\flat(e_j)$ where $\{e_1, e_2, ..., e_{2(n-1)}\}$ is any basis of $\mathcal{E}$ and $[m^{ij}]$ is a matrix inverse to $[m_{ij}]$, where $m_{ij} = m(e_i, e_j)$. Note that if $f \in C^\infty(M)$ then $div_{\mathcal{E}}(fX) = f div_{\mathcal{E}} X$ in the case $X \in \Gamma(\mathcal{D})$. Let $\xi \in \Gamma(\mathcal{D})$ be a unit local section of $\mathcal{D}$. Then $\{\xi, J\xi\}$ is an orthonormal basis of $\mathcal{D}$. Let $\theta(X) = g(\xi, X)$ and $J\theta = -\theta \circ J$ which means that $J\theta(X) = g(J\xi, X)$. Let us denote by κ the function

$$\kappa = \sqrt{(div_{\mathcal{E}}\xi)^2 + (div_{\mathcal{E}} J\xi)^2}. \tag{1.1}$$

The function κ does not depend on the choice of a section ξ. In fact, if $\xi' = a\xi + bJ\xi$, where $a, b \in C^\infty(dom\xi)$ and $a^2 + b^2 = 1$ is another unit section of $\mathcal{D}$, then $J\xi' = -b\xi + aJ\xi$ and

$$\begin{aligned}(div_{\mathcal{E}}\xi')^2 + (div_{\mathcal{E}}J\xi')^2 &= (a div_{\mathcal{E}}\xi + b div_{\mathcal{E}}J\xi)^2 + (-b div_{\mathcal{E}}\xi + a div_{\mathcal{E}}J\xi)^2 \\ &= (div_{\mathcal{E}}\xi)^2 + (div_{\mathcal{E}}J\xi)^2.\end{aligned} \tag{1.2}$$

Hence κ is a well defined, continuous function on M, which is smooth in the open set $U = \{x : \kappa(x) \neq 0\}$. We shall now show that on U there is a smooth, global unit section $\xi \in \Gamma(U, \mathcal{D})$ defined uniquely up to a sign such that $div_{\mathcal{E}} J\xi = 0$. Namely, if ξ' is a local unit section of $\Gamma(U, \mathcal{D})$ then then the section $\xi = \frac{1}{\kappa}((div_{\mathcal{E}}\xi')\xi' + (div_{\mathcal{E}}J\xi)J\xi')$ satisfies $div_{\mathcal{E}}J\xi = \frac{1}{\kappa}((div_{\mathcal{E}}\xi')(div_{\mathcal{E}}J\xi') - (div_{\mathcal{E}}J\xi')(div_{\mathcal{E}}\xi')) = 0$ and does not depend on the choice of ξ'. On the other hand it is clear that the only other such smooth section is $-\xi$. The section ξ constructed above and defined on $U \subset M$ we shall call the principal section of $\mathcal{D}$. Note that $div_{\mathcal{E}}\xi = \kappa$.

2. Curvature tensor of a QCH Kähler manifold. We shall recall some results from [G-M-1]. Let $R(X,Y)Z = ([\nabla_X, \nabla_Y] - \nabla_{[X,Y]})Z$ and let us write

$$R(X,Y,Z,W) = g(R(X,Y)Z,W).$$

If R is the curvature tensor of a QCH Kähler manifold (M,g,J), then there exist functions $a, b, c \in C^\infty(M)$ such that

$$R = a\Pi + b\Phi + c\Psi, \tag{2.1}$$

where Π is the standard Kähler tensor of constant holomorphic curvature i.e.

$$\begin{aligned}\Pi(X,Y,Z,U) = \frac{1}{4}(&g(Y,Z)g(X,U) - g(X,Z)g(Y,U) \\ &+g(JY,Z)g(JX,U) - g(JX,Z)g(JY,U) - 2g(JX,Y)g(JZ,U)),\end{aligned} \tag{2.2}$$

the tensor Φ is defined by the following relation

$$\begin{gathered}\Phi(X,Y,Z,U)=\frac{1}{8}(g(Y,Z)h(X,U)-g(X,Z)h(Y,U)\\ +g(X,U)h(Y,Z)-g(Y,U)h(X,Z)+g(JY,Z)\omega(X,U)\\ -g(JX,Z)\omega(Y,U)+g(JX,U)\omega(Y,Z)-g(JY,U)\omega(X,Z)\\ -2g(JX,Y)\omega(Z,U)-2g(JZ,U)\omega(X,Y)),\end{gathered} \tag{2.3}$$

and finally

$$\Psi(X,Y,Z,U)=-\omega(X,Y)\omega(Z,U)=-(\omega\otimes\omega)(X,Y,Z,U). \tag{2.4}$$

Let $V=(V,g,J)$ be a real $2n$ dimensional vector space with complex structure J which is skew-symmetric with respect to the scalar product g on V. Let assume further that $V=D\oplus E$ where D is a 2-dimensional, J-invariant subspace of V, E denotes its orthogonal complement in V. Note that the tensors Π,Φ,Ψ given above are of Kähler type. It is easy to check that for a unit vector $X\in V$ $\Pi(X,JX,JX,X)=1,\Phi(X,JX,JX,X)=|X_D|^2,\Psi(X,JX,JX,X)=|X_D|^4$, where X_D means an orthogonal projection of a vector X on the subspace D and $|X|=\sqrt{g(X,X)}$. It follows that for a tensor (2.1) defined on V we have

$$R(X,JX,JX,X)=\phi(|X_D|)$$

where $\phi(t)=a+bt^2+ct^4$.

Now let us assume that (M,g,J) is a QCH Kähler manifold of dimension $2n\geq 6$ and let X be a local unit section of $\mathcal{D}$ and $\eta(Z)=g(X,Z)$. Let us define two 1-forms ϵ,ϵ^* by the formulas: $\epsilon(Z)=g(p_{\mathcal{E}}(\nabla_X X),Z)=g(\nabla_X X,Z)-pJ\eta(Z)$, $\epsilon^*(Z)=g(p_{\mathcal{E}}(\nabla_{JX}JX),Z)=g(\nabla_{JX}JX,Z)-p^*\eta(Z)$ where $p=g(\nabla_X X,JX)$,$p^*=g(\nabla_{JX}JX,X)$ and $p_{\mathcal{E}}$ denotes the orthogonal projection on $\mathcal{E}$. Note that the distribution $\mathcal{D}$ is integrable if and only if $\epsilon+\epsilon^*=0$ (see [G-M-1], Lemma 3.3). In fact for $Z\in\Gamma(\mathcal{E})$ we have

$$\begin{gathered}g([X,JX],Z)=g(\nabla_X JX-\nabla_{JX}X,Z)=g(J\nabla_X JX-J\nabla_{JX}X,JZ)\\ =-g(\nabla_X X+\nabla_{JX}JX,JZ)=-(\epsilon(JZ)+\epsilon^*(JZ)).\end{gathered}$$

Let $\{Z_\lambda\}$ be any complex basis of the complex subbundle $\mathcal{E}^{1,0}$ of the complex tangent bundle $T^cM=\mathbb{C}\otimes TM$. We also write $Z_{\bar\lambda}=\bar{Z_\lambda}$. Then the Bianchi identity for the tensor R of the form (2.1) gives the following relations (see Theorem 3.5 in [G-M-1]) :

$$\begin{gathered}\nabla a=\frac{bdiv_{\mathcal{E}}X}{2(n-1)}X+\frac{bdiv_{\mathcal{E}}JX}{2(n-1)}JX,\\ \nabla b=\frac{(b+4c)div_{\mathcal{E}}X}{n-1}X+\frac{(b+4c)div_{\mathcal{E}}JX}{n-1}JX,\\ b\nabla_{Z_\lambda}\eta(Z_\mu)=0,c\nabla_{Z_\lambda}\eta(Z_\mu)=0,\\ b(\nabla_{Z_\lambda}\eta(Z_{\bar\mu})-\frac{div_{\mathcal{E}}X}{2(n-1)}g_{\lambda\bar\mu}+\frac{div_{\mathcal{E}}JX}{2(n-1)}\Omega_{\lambda\bar\mu})=0\\ c(\nabla_{Z_\lambda}\eta(Z_{\bar\mu})-\frac{div_{\mathcal{E}}X}{2(n-1)}g_{\lambda\bar\mu}+\frac{div_{\mathcal{E}}JX}{2(n-1)}\Omega_{\lambda\bar\mu})=0\\ b\epsilon(Z_\lambda)=0,\ b\epsilon^*(Z_\lambda)=0\\ c(\epsilon(Z_\lambda)+\epsilon^*(Z_\lambda))=dc(Z_\lambda).\end{gathered} \tag{2.5}$$

In particular in $U=\{x\in M:\kappa(x)\neq 0\}$ we obtain

$$\nabla a=\frac{b}{2(n-1)}\kappa\xi \qquad \nabla b=\frac{(b+4c)}{n-1}\kappa\xi, \tag{2.6}$$

where ξ is the principal section of the bundle $\mathcal{D}$. In the rest of the paper we shall assume that the set $U=\{x:\kappa(x)\neq 0\}$ is non-empty and the set $B=\{x\in U: b(x)=0\}$ has an empty interior. Thus we obtain

$$\begin{gathered}\nabla_{Z_\lambda}\eta(Z_\mu)=0,\\ \nabla_{Z_\lambda}\eta(Z_{\bar\mu})-\frac{div_{\mathcal{E}}X}{2(n-1)}g_{\lambda\bar\mu}+\frac{div_{\mathcal{E}}JX}{2(n-1)}\Omega_{\lambda\bar\mu}=0\\ \epsilon=0,\ \epsilon^*=0.\end{gathered} \tag{2.7}$$

It follows that the distribution $\mathcal{D}_U$ is integrable. In U we get

$$\nabla_{Z_\lambda}\eta(Z_{\bar\mu})=\frac{\kappa}{2(n-1)}g_{\lambda\bar\mu}. \tag{2.8}$$

Let ξ be the principal section of $\mathcal{D}$ defined in U and let $\theta=\xi^\flat$ be the dual 1-form of ξ. Then in U

$$da=\frac{b}{2(n-1)}\kappa\theta, db=\frac{(b+4c)}{n-1}\kappa\theta. \tag{2.9}$$

It follows that the distribution $\Delta=\{X\in TU:\theta(X)=0\}$ defined in U is integrable. From (2.7) and (2.8) it follows that $\nabla\theta(JX,JY)=\nabla\theta(X,Y)$ for $X,Y\in\Gamma(\mathcal{E})$. It is also clear that

$$dJ\theta_{|\mathcal{E}}=\frac{\kappa}{n-1}\Omega_{|\mathcal{E}}.$$

Thus the distribution $\mathcal{E}_{|U}$ is the so called B_0 -distribution defined in [G-M-1]. Consequently we obtain (see [G-M-1], Lemma 5.1):

$$\nabla_X\theta(Y)=\frac{\kappa}{2(n-1)}m(X,Y)+p\theta(x)J\theta(Y)-p^*J\theta(X)J\theta(Y), \tag{2.10}$$

If dim $M=2n\geq 6$ we also have $d\theta=p\,\theta\wedge J\theta$, $p=g(\nabla_\xi\xi,J\xi)=0$ and consequently (see Lemma 5.2 in [G-M-1] and its proof): $\nabla_\xi\xi=0, d\theta=0$,

$$d\ln\kappa=-(\frac{\kappa}{n-1}+p^*)\theta, dp^*\wedge\theta=0. \tag{2.11}$$

Thus

$$\nabla_X\theta(Y)=\frac{\kappa}{2(n-1)}m(X,Y)-p^*J\theta(X)J\theta(Y), \tag{2.12}$$

If (M,g,J) is a QCH Kähler manifold then one can show that the Ricci tensor ρ of (M,g,J) satisfies the equation

$$\rho(X,Y)=\lambda m(X,Y)+\mu h(X,Y) \tag{2.13}$$

where $\lambda=\frac{n+1}{2}a+\frac{b}{4}, \mu=\frac{n+1}{2}a+\frac{n+3}{4}b+c$ are eigenvalues of ρ (see [G-M-1], Corollary 2.1 and Remark 2.1.) In particular the distributions $\mathcal{E},\mathcal{D}$ are eigendistributions of the tensor ρ corresponding to the eigenvalues λ,μ of ρ.

3. Local holomorphic Killing vector field on U. Let (M,g,J) be a QCH Kähler manifold of dimension $2n\geq 6$. We shall show in this section that for every $x\in U$ there exists an open neighborhood $V\subset U$ of x and a function $f\in C^\infty(V)$ such that $X_V=fJ\xi$ is a Killing vector field in V, which we shall call a local special Killing vector field. Let V be a geodesically convex neighborhood of x in U. Then V is contractible. Note that the form $\phi=-p^*\theta$, where $p^*=g(\nabla_{J\xi}J\xi,\xi)$ is closed in U, since by (2.11), $d\phi=-dp^*\wedge\theta=0$. Consequently there exists a function $F\in C^\infty(V)$ such that

$$dF=\phi=-p^*\theta. \tag{3.1}$$

Let $f=\exp\circ F$. From (2.12) it follows that

$$\nabla_X J\theta(Y)=\frac{\kappa}{2(n-1)}\Omega_m(X,Y)+p^*J\theta(X)\theta(Y), \tag{3.2}$$

Now let $\psi=(fJ\xi)^\flat=fJ\theta$. We shall show that $\nabla_X\psi(Y)=-\nabla_Y\psi(X)$ which means that the field $fJ\xi$ is a Killing vector field in V.We get

$$\nabla_X(fJ\theta)(Y)=XfJ\theta(Y)+f\frac{\kappa}{2(n-1)}\Omega_m(X,Y)+fp^*J\theta(X)\theta(Y), \tag{3.3}$$

Since $Xf=fXF=-fp^*\theta(X)$ we obtain

$$\begin{aligned}\nabla_X(fJ\theta)(Y)=f\frac{\kappa}{2(n-1)}\Omega_m(X,Y)-fp^*\theta\wedge J\theta(X,Y)=\\ f\frac{\kappa}{2(n-1)}\Omega_m(X,Y)-fp^*\omega(X,Y),\end{aligned} \tag{3.4}$$

which proves our claim. Note that if F_1 is another solution of (3.1) then $F_1=F+D$ for a certain constant $D\in\mathbb{R}$. It follows that $f_1=\exp F_1=CF$, where $C=\exp D$. Consequently $X_1=f_1J\xi=CfJ\xi=CX$. Recall here the well known general fact, that if $X,Y\in\mathfrak{iso}(M)$ are Killing vector fields on connected Riemannian manifold M, $X\neq 0$ and $Y=fX$ for a certain $f\in C^\infty(M)$ then f is constant.

Let $\phi=f\xi^\flat$. Now it is clear that

$$\nabla_X\phi(Y)=\nabla_X(f\theta)(Y)=f\frac{\kappa}{2(n-1)}m(X,Y)-fp^*h(X,Y) \tag{3.5}$$

and $\nabla_X(f\theta)(Y)=\nabla_Y(f\theta)(X)$. Consequently $d(f\theta)=0$. It follows that there exists a function $\tau\in C^\infty(V)$ such that $f\xi=\nabla\tau$. Consequently $X_V=fJ\xi=$

$J(\nabla\tau)$, which means that X_V is a holomorphic Killing vector field with a Kähler potential τ.

4. Special Jacobi fields along geodesics in U. Let $c : [0,l] \to M$ be a unit geodesic such that $c([0,l)) \subset U$ and $c(l) \in K = \{x \in M : \kappa(x) = 0\}$. A vector field C, which is a Jacobi field along c i.e. $\nabla^2_{\dot c}C - R(\dot c, C)\dot c = 0$, will be called a special Jacobi field if there exists an open, geodesically convex neighborhood V of $c(0)$ such that $C(0) = X_V(c(0)), \nabla_{\dot c}C(0) = \nabla_{\dot c}X_V(c(0))$. If $im\ c \cap V = c([0,\epsilon))$ then it follows that $X_V(c(t)) = C(t)$ for all $t \in [0,\epsilon)$. We have the following lemma:

Lemma 4.1. *Let us assume that a vector field C along a geodesic c is a special Jacobi field along c. Then*

$$\lim_{t\to l}|C(t)| = 0,$$

and

$$g(\dot c, C) = 0.$$

Proof. Let us note that $g(\nabla_{\dot c}C, \dot c) = 0$ since this property is valid for Killing vector fields. It follows that the function $g(\dot c, C)$ is constant. Let $k \in (0,l)$. Then $c([0,k]) \subset U$. For every $t \in [0,k)$ there exists a geodesically convex open neighborhood V_t of the point $c(t)$ and a special Killing vector field $X_{V_t} = f_tJ\xi$ on V_t defined in Section 3. The field X_{V_t} is defined uniquely up to a constant factor. From the cover $\{V_t\} : t \in [0,k]$ of the compact set $c([0,k])$ we can choose a finite subcover $\{V_{t_1}, V_{t_2}, ..., V_{t_m}\}$. Let c_i be the part of geodesic c contained in $V_i = V_{t_i}$, i.e. $im\ c \cap V_i = im\ c_i = c((t_i, t_{i+1}))$. We define the Killing vector field X_i in every $V_i = V_{t_i}$ by induction in such a way that $X_1 = X_V$ on $V_1 \cap V$ and $X_i = X_{i+1}$ on $V_i \cap V_{i+1}$. Let $X_i = f_iJ\xi$. Note that $C(t) = X_i \circ c(t) = f_iJ\xi \circ c(t)$ for $t \in (t_i, t_{i+1})$. Consequently, on V_i, $|C| = f_i$. From (2.11) and (3.1) it follows that

$$d\ln\kappa = d\ln f_i - \frac{\kappa}{n-1}\theta.$$

Hence

$$\frac{d}{dt}\ln\kappa\circ c(t) = \frac{d}{dt}\ln|C(t)| - \frac{\kappa}{n-1}\theta(\dot c(t)), \tag{4.1}$$

and

$$\frac{d}{dt}\ln\frac{\kappa\circ c(t)}{|C(t)|} = -\frac{\kappa}{n-1}\theta(\dot c(t)). \tag{4.2}$$

Consequently

$$\ln\frac{\kappa\circ c(k)}{|C(k)|} - \ln\frac{\kappa\circ c(0)}{|C(0)|} = -\frac{1}{n-1}\int_0^k \kappa\theta(\dot c(t))dt. \tag{4.3}$$

Hence

$$\ln|C(k)| = \ln\kappa\circ c(k) - \ln\frac{\kappa\circ c(0)}{|C(0)|} + \frac{1}{n-1}\int_0^k \kappa\theta(\dot c(t))dt. \tag{4.4}$$

Note that

$$|\int_0^k \kappa\theta(\dot{c}(t))dt| \leq \int_0^k |\kappa\theta(\dot{c}(t))|dt \leq \int_0^k \kappa|\dot{c}(t)|dt \leq \int_0^k \kappa dt. \tag{4.5}$$

Let $\kappa_0 = sup\{\kappa(x) : x \in c([0,l])\}$. From (4.4) it follows that

$$\ln|C(k)| \leq \ln\kappa\circ c(k) - \ln\frac{\kappa\circ c(0)}{|C(0)|} + \frac{1}{n-1}\kappa_0 l. \tag{4.6}$$

Consequently

$$\limsup_{k\to l-}\ln|C(k)| \leq \lim_{k\to l-}\ln\kappa\circ c(k) - \ln\frac{\kappa\circ c(0)}{|C(0)|} + \frac{1}{n-1}\kappa_0 l = -\infty. \tag{4.7}$$

From (4.7) it is clear that $\lim_{k\to l-}|C(k)| = 0$. Since $|g(\dot{c}(t), C(t)) \leq |C(t)|$ and $g(\dot{c}, C)$ is constant it follows that $|g(\dot{c}(t), C(t))| \leq \lim_{t\to l}|C(t)| = 0$ which means that $g(\dot{c}, C) = 0$.♢

5. Global holomorphic Killing vector field on M. From now on we assume that (M, g, J) is a complete QCH Kähler manifold, $\dim M \geq 6$ and the set $U = \{x \in M : \kappa(x) \neq 0\}$ is non-empty and B has an empty interior. Let $K = \{x \in M : \kappa(x) = 0\}$.

Theorem 5.1. *The set U is connected and the set K has an empty interior.*

Proof. Let U_1 be a non-empty component of the set $U = M - K$. Let $x_0 \in U_1$ and let us assume the set $int\, K \cup (U - U_1)$ is non-empty. Let $x_1 \in int\, K \cup (U - U_1)$. Then $x_1 = \exp_{x_0} lX$ for a certain unit vector $X \in T_{x_0}M$ and $l > 0$. Let $V \subset U_1$ be a geodesically convex neighborhood of x_0 and $X_V = f_V J\xi$ the local Killing vector field on V. Let C be the Jacobi vector field along the geodesic $c(t) = \exp_{x_0} tX$ satisfying the initial conditions:$C(0) = X_V(x_0), \nabla_{\dot{c}}C(0) = \nabla_{\dot{c}}X_V(x_0)$. It follows that the field C is a special Jacobi field along c. In particular $g(X, X_V(x_0)) = 0$ since the geodesic c meets K. Since the set $int\, K \cup (U - U_1)$ is open there exists an open neighborhood $W \subset int\, K \cup (U - U_1)$ of the point x_1. The mapping $T_{x_0}M \ni Y \to \exp_{x_0} lY \in M$ is continuous hence there exists an open neighborhood P of X in $T_{x_0}M$ such that $\exp_{x_0} lP \subset int\, K \cup (U - U_1)$. We can find a vector $X_1 \in P$ such that $g(X_1, X_V(x_0)) \neq 0$. The field C_1 along the geodesic $d(t) = \exp_{x_0} tX_1$ defined by the initial conditions $C_1(0) = X_V(x_0), \nabla_{\dot{d}}C_1(0) = \nabla_{\dot{d}(0)}X_V$ is a special Jacobi field along a geodesic d which meets K. It follows that $g(\dot{d}, C_1) = 0$ along d. In particular for $t = 0$ we obtain $g(X_1, X_V(x_0)) = 0$ which is a contradiction. Consequently $int\, K \cup (U - U_1) = \emptyset$.♢

Corollary 5.2. Let us assume that (M, g, J) is a complete QCH Kähler manifold, $\dim M \geq 6$, the set $U = \{x \in M : \kappa(x) \neq 0\}$ is non-empty and B has an empty interior. Then the distribution $\mathcal{D}$ is totally geodesic, i.e. $\nabla_X Y \in \Gamma(\mathcal{D})$ for every $X, Y \in \Gamma(\mathcal{D})$.

Proof. Let $\nabla_X Y = p_{\mathcal{D}}(\nabla_X Y) + \alpha(X, Y)$ where $\alpha(X, Y) = p_{\mathcal{E}}(\nabla_X Y)$ and α is a section of the bundle $\mathcal{D}^* \otimes \mathcal{D}^* \otimes \mathcal{E}$. Since $\mathcal{D}$ is integrable α is a symmetric two-form. We shall show that $\alpha = 0$. The bundle $\mathcal{D}_{|U}$ is spanned by the sections $\xi, J\xi$ and hence locally by $-f\xi = JX, X = fJ\xi$ where X is a special Killing field defined in

$dom f = V \subset U$. Note that $g(X,X) = f^2$ and $\nabla_X X = -\frac{1}{2}\nabla[g(X,X)] = -f\nabla f$. Thus (see (3.1)) $\nabla_X X = f^2 p^*\xi \in \Gamma(\mathcal{D})$. Since X is a holomorphic Killing vector field it follows that $[X, JX] = L_X(JX) = J[X,X] = 0$. Hence $\nabla_X JX = \nabla_{JX} X$ and $\nabla_X JX = J\nabla_X X = Xf\xi + f^2\nabla_\xi\xi = Xf\xi \in \Gamma(\mathcal{D})$ since $\nabla_\xi\xi = 0$. Finally $\nabla_{JX} JX = J(\nabla_{JX} X) = XfJ\xi \in \Gamma(\mathcal{D})$. It follows that $\alpha(X,X) = \alpha(X,JX) = \alpha(JX,JX) = 0$ and $\alpha_{|U} = 0$. Since U is dense in M it follows that $\alpha = 0$ in M. $\diamondsuit$

Let $x_0 \in U$ and let $V = \exp_{x_0} W$ be a geodesically convex open neighborhood of x_0, where $W \subset T_{x_0}M$ is a star shaped open neighborhood of $0 \in T_{x_0}M$. For every $X \in W$ let $l(X) = \sup\{t : tX \in W\}$. Hence if the sphere $S_\epsilon = \{X \in T_{x_0}M : |X| = \epsilon\}$ is contained in $W - \exp_{x_0}^{-1}(K)$ then $W = \{tX : X \in S_\epsilon, t \in [0, l(X))\}$. In an open neighborhood $V' \subset V - K$ of x_0 there is defined a special Killing vector field $X_{V'} = f_{V'}J\xi$. Let $Z = X_{V'}(x_0) \in T_{x_0}M$ so that $H = \{X \in T_{x_0}M : g_{x_0}(X,Z) = 0\}$ is a hyperplane in $T_{x_0}M$. Note that $\exp_{x_0}^{-1}(V \cap K) \subset H$. Let a function $k : S_\epsilon \to \mathbb{R}$ be defined as follows: $k(X) = l(X)$ if $x \notin H$ and $k(X) = \inf\{t > 0 : \exp(tX) \in K\}$ if $X \in H$. The set $W' = \{tX : X \in S_\epsilon, t \in [0, k(X))\}$ is open and star shaped. Note that $V'' = \exp_{x_0} W' \subset V - K \subset U$, V'' is dense in V, and V'' is contractible. It follows that in V'' there is defined a special local Killing vector field $X_{V''}$. We can assume that $X_{V'} = X_{V''}$ on V'. Now we can prove:

Lemma 5.3 *On every geodesically convex open set V in M can be defined a holomorphic Killing vector field X such that for every open geodesically convex set $W \subset V \cap U$ the restriction $X_{|W}$ is a special Killing vector field on W.*

Proof. We shall use the notation introduced above. Since V'' is contractible there exists a special Killing vector field $X_{V''}$ defined on V''. Let us define a differentiable field X on V by the formula: $X(\exp_{x_0} u) = J_u(1)$ where $u \in W$ and J_u is a Jacobi vector field along a geodesic $c(t) = \exp_{x_0}(tu)$ satisfying the initial conditions:$J_u(0) = X_{V''}(x_0), \nabla_{\dot c} J_u(0) = \nabla_u X_{V''}$. It is clear that X is a differentiable vector field and that $X_{|V''} = X_{V''}$. Since the set V'' is dense in V it follows that X is a Killing vector field in V. In fact if we write $T = \nabla X$ then for every $Y, Z \in \mathfrak{X}(V)$ we have $g(TY, Z) = -g(Y, TZ)$ on V'' and both sides are differentiable functions on V. Thus the relation remains valid on V which means that X is a Killing vector field on V. Note that $X = 0$ on $V \cap K$. Since equation (3.5) is valid on the open, dense subset $V - K$ of V it follows that the form $\phi = -(JX)^\flat$ satisfies the relation $\nabla_Y\phi(Z) = \nabla_Z\phi(Y)$ for every $Y, Z \in \mathfrak{X}(V)$. Consequently $JX = -\nabla\tau$ for a certain function $\tau \in C^\infty(V)$ and $X = J\nabla\tau$ on V. Since $\nabla_{JY}\phi(JZ) = \nabla_Y\phi(Z)$ on V'' and hence on V for every $Y, Z \in \mathfrak{X}(V)$ it is clear that X is holomorphic. $\diamondsuit$

Let V_1, V_2 be two open, geodesically convex sets. Then the set $V_1 \cap V_2 - K$ is connected. The proof of this is similar to the proof of Theorem 5.1. Since the sets $V_1'' \subset V_1 - K$, $V_2'' \subset V_2 - K$ are contractible there exist special Killing vector fields $X_1 = f_1J\xi, X_2 = f_2J\xi$ on these sets. These Killing fields can be extended on $V_1 - K, V_2 - K$ in such a way that $f_i = \exp F_i$ where F_i satisfy equation (3.1). Consequently $d(F_1 - F_2) = 0$ on the connected set $V_1 \cap V_2 - K$. Hence $F_1 = F_2 + D$ in $V_1 \cap V_2 - K$ for a constant $D \in \mathbb{R}$. Consequently $X_1 = C_{12}X_2$ where $C_{12} = \exp D$. The fields X_1, X_2 can be extended to the Killing fields on V_1, V_2 respectively such that $X_{i|K\cap V_i} = 0$. It is clear that for these extensions which we also denote by X_1, X_2 the equation

$$X_1 = C_{12}X_2$$

holds on $V_1 \cap V_2$.

Let us recall the definition of a special Kähler-Ricci potential ([D-M-1],[D-M-2]).

Definition. *A nonconstant function $\tau \in C^\infty(M)$, where (M, g, J) is a Kähler manifold, is called a special Kähler-Ricci potential if the field $X = J(\nabla\tau)$ is a Killing vector field and, at every point with $d\tau \neq 0$ all nonzero tangent vectors orthogonal to the fields X, JX are eigenvectors of both $\nabla d\tau$ and the Ricci tensor ρ of (M, g, J).*

Now we shall prove:

Theorem 5.4. *Let (M, g, J) be a complete QCH Kähler manifold of dimension $2n \geq 6$. Let $intB = \emptyset$ and $U \neq \emptyset$. If $H^1(M, \mathbb{R}) = 0$ then there exists on M a non-zero holomorphic Killing vector field $X = J(\nabla\tau)$ with a special Kähler-Ricci potential τ.*

Proof. Let $\{V_i\} : i \in I$ be a cover of M by geodesically convex, open sets V_i. Let X_i be a Killing vector field on V_i constructed in Lemma 5.2. Let $V_i \cap V_j \neq \emptyset$. Then there exist constants $C_{ij} > 0$ such that $X_i = C_{ij} X_j$ on $V_i \cap V_j$. These constants satisfy the co-cycle condition $C_{ij} C_{jk} C_{ki} = 1$. Consequently the constants $D_{ij} = \ln C_{ij} \in \mathbb{R}$ satisfy the co-cycle condition $D_{ij} + D_{jk} + D_{ki} = 0$. It follows that $\{D_{ij}\}$ is a co-cycle in the first Čech cohomology group $\check{H}^1(\{V_i\}, \mathbb{R})$. Since $\{V_i\}$ is a good cover of M it follows that $\check{H}^1(\{V_i\}, \mathbb{R}) = \check{H}^1(M, \mathbb{R}) = 0$. Consequently there exists a co-cycle $\{D_i\} \in Z^0(\{V_i\}, \mathbb{R})$ such that $\{D_{ij}\} = \delta(\{D_i\})$. This means that $D_{ij} = D_j - D_i$. Let $C_i = \exp D_i$. Then $C_{ij} = \frac{C_j}{C_i}$. Let us define the field X on M by the formula $X_{|V_i} = C_i X_i$. Then it is clear that X is a well defined, global vector field and $X \in \mathfrak{X}(M)$. Since $X_{|V_i}$ is a Killing vector field on every V_i it follows that X is a Killing vector field. Now let $\phi = -(JX)^\flat$. Then $d\phi = 0$, since this equation is satisfied on every V_i. On the other hand the first de Rham group of M vanishes: $H^1(M, \mathbb{R}) = \check{H}^1(M, \mathbb{R}) = 0$. Consequently there exists a function $\tau \in C^\infty(M)$, such that $\phi = d\tau$. Note also that $\nabla d\phi$ is Hermitian, which means that X is holomorphic. Thus $X = J(\nabla\tau)$ is a holomorphic Killing vector field with a Killing potential τ. Note that in view of (2.12) and (3.5) the special Killing field constructed by us is a Killing vector field with a special Kähler-Ricci potential τ. $\diamondsuit$

Corollary 5.5. *Let (M, g, J) be a complete QCH Kähler manifold of dimension $2n \geq 6$. Let $intB = \emptyset$, $U \neq \emptyset$ and let $(\tilde{M}, \tilde{g})$ be the Riemannian universal covering space of (M, g, J). Then there exists on $(\tilde{M}, \tilde{g})$ a non-zero holomorphic Killing vector field with a special Kähler-Ricci potential.*

6. Construction of QCH Kähler manifolds.

In our construction we shall follow Bérard Bergery (see [Ber], [S]) rather then Derdziński and Maschler, although we shall use the classification theorem by Derdziński and Maschler ([D-M-1]) to classify QCH Kähler manifolds. Note that these two approaches are equivalent (see [D-M-2]). Let (N, h, J) be a simply connected Kähler Einstein manifold, which is not Ricci flat and $\dim N = 2m > 2$. Let $s \geq 0, L > 0, s \in \mathbb{Q}, L \in \mathbb{R}$ and $r : [0, L] \to \mathbb{R}$ be a positive, smooth function on $[0, L]$ with $r'(t) > 0$ for $t \in (0, L)$, which is even at 0 and L, i.e. there exists an $\epsilon > 0$ and even, smooth functions $r_1, r_2 : (-\epsilon, \epsilon) \to \mathbb{R}$ such that $r(t) = r_1(t)$ for $t \in [0, \epsilon)$ and $r(t) = r_2(L - t)$ for $t \in (L-\epsilon, L]$. If $s \neq 0$ then it is clear that the function $f = \frac{2}{s}rr'$ is positive on $(0, L)$ and $f(0) = f(L) = 0$. Let P be a circle bundle over N classified by the integral cohomology class $\frac{s}{2}c_1(N) \in H^2(N, \mathbb{R})$ if $c_1(M) \neq 0$. Let q be the unique positive integer such that $c_1(N) = q\alpha$ where $\alpha \in H^2(N, \mathbb{R})$ is an indivisible integral class. Then

$$s = \frac{2k}{q}; k \in \mathbb{Z}.$$

It is known that $q = n$ if $N = \mathbb{CP}^{n-1}$ (see [Bes], p.273). Note that $c_1(N) = \{\frac{1}{2\pi}\rho_N\} = \{\frac{\tau_N}{4m\pi}\Omega_N\}$ where $\rho_N = \frac{\tau_N}{2m}\Omega_N$ is the Ricci form of (N, h, J), τ_N is the scalar curvature of (N, h) and Ω_N is the Kähler form of (N, h, J). We can assume that $\tau_N = \pm 4m$. In the case $c_1(N) = 0$ we shall assume that (N, h, J) is a Hodge manifold, i.e. the cohomology class $\{\frac{s}{2\pi}\Omega_N\}$ is an integral class. On the bundle $p : P \to N$ there exists a connection form θ such that $d\theta = sp^*\Omega_N$ where $p : P \to N$ is the bundle projection. Let us consider the manifold $(0, L) \times P$ with the metric

$$g = dt^2 + f(t)^2\theta^2 + r(t)^2 p^*h, \tag{6.1}$$

if $s \neq 0$ and the metric

$$g = dt^2 + f(t)^2\theta^2 + p^*h, \tag{6.2}$$

if $s = 0$. The metric (6.2) is a Kähler product metric. The metric 6.1 is Kähler if and only if $f = \frac{2rr'}{s}$. We shall prove it in section 8. It is known that the metric (6.1) extends to a metric on a sphere bundle $M = P \times_{S^1} \mathbb{CP}^1$ if and only if the function r is positive and smooth on $(0, L)$, even at the points $0, L$, the function f is positive, smooth an odd at the points $0, L$ and additionally

$$f'(0) = 1, f'(L) = -1. \tag{6.3}$$

If $f = \frac{2rr'}{s}$ for r as above, then (6.3) means that

$$2r(0)r''(0) = s, 2r(L)r''(L) = -s. \tag{6.4}$$

7. Circle bundles. Let (N, h, J) be a Kähler manifold with integral class $\{\frac{s}{2\pi}\Omega_N\}$, where $s \in \mathbb{Q}$ and let $p : P \to N$ be a circle bundle with a connection form θ such that $d\theta = s\Omega_N$ (see [K]). We shall assume that (N, h) is Einstein if *dim* $N > 2$. Let us consider a Riemannian metric g on P given by

$$g = a^2\theta \otimes \theta + b^2 p^*h \tag{7.1}$$

where $a, b \in \mathbb{R}$. Let ξ be the fundamental vector field of the action of S^1 on P i.e. $\theta(\xi) = 1, L_\xi g = 0$. It follows that $\xi \in \mathfrak{iso}(P)$ and $a^2\theta = g(\xi, .)$. Consequently

$$a^2 d\theta(X, Y) = 2g(TX, Y) \tag{7.2}$$

for every $X, Y \in \mathfrak{X}(P)$ where $TX = \nabla_X\xi$. Note that $g(\xi, \xi) = a^2$ is constant, hence $T\xi = 0$. On the other hand $d\theta(X, Y) = sp^*\Omega_N(X, Y) = sh(Jp(X), p(Y))$. Note that there exists a tensor field $\tilde{J}$ on P such that $\tilde{J}\xi = 0$ and $\tilde{J}(X) = (JX_*)^H$ where $X = X_*^H \in TP$ is the horizontal lift of $X_* \in TN$ (i.e. $\theta(X_*^H) = 0$) and $X_* = p(X)$. Indeed $L_\xi T = 0$ and $T\xi = 0$ hence T is the horizontal lift of the tensor $\tilde{T}$. Now $\tilde{J} = \frac{2b^2}{sa^2}\tilde{T}$. Since $T\xi = 0$ we get $\nabla T(X,\xi) + T^2X = 0$ and $R(X,\xi)\xi = -T^2X$. Thus $g(R(X,\xi)\xi, X) = ||TX||^2$ and

$$\rho(\xi,\xi) = ||T||^2 = \frac{sa^4}{4b^4}2m.$$

Consequently

$$\lambda = \rho(\frac{\xi}{a},\frac{\xi}{a}) = \frac{1}{a^2}||T||^2 = \frac{s^2a^2}{4b^4}m. \tag{7.3}$$

We shall compute the O'Neill tensor A (see [ON]) of the Riemannian submersion $p : (P,g) \to (N, b^2h)$. We have

$$A_EF = \mathcal{V}(\nabla_{\mathcal{H}E}\mathcal{H}F) + \mathcal{H}(\nabla_{\mathcal{H}E}\mathcal{V}F).$$

Let us write $u = \mathcal{V}(\nabla_{\mathcal{H}E}\mathcal{H}F)$ and $v = \mathcal{H}(\nabla_{\mathcal{H}E}\mathcal{V}F)$. The vertical component of a field E equals $\theta(E)\xi$. If $X, Y \in \mathcal{H}$ then

$$g(\nabla_XY, \frac{1}{a}\xi) = \frac{1}{a}(Xg(Y,\xi) - g(Y, \nabla_X\xi)) = -\frac{1}{a}g(TX,Y) = \frac{1}{a}g(X,TY). \tag{7.4}$$

Hence $u = \frac{1}{a}g(E - \theta(E)\xi, T(F - \theta(F)\xi)\frac{\xi}{a} = \frac{1}{a^2}g(E,TF)\xi$. Note that $\mathcal{H}(\nabla_X f\xi) = f\mathcal{H}(\nabla_X\xi) = fTX$, hence

$$v = \mathcal{H}(\nabla_{\mathcal{H}E}\mathcal{V}F) = \theta(F)T(E) = \frac{1}{a^2}g(\xi,F)TE.$$

Consequently

$$A_EF = \frac{1}{a^2}(g(E,TF)\xi + g(\xi,F)TE). \tag{7.5}$$

If $U, V \in \mathcal{H}$ then

$$||A_UV||^2 = \frac{1}{a^2}g(E,TF)^2 = \frac{s^2a^2}{4b^4}g(E,\tilde{J}F)^2.$$

If E is horizontal and F is vertical then

$$A_EF = \frac{1}{a^2}g(\xi,F)TE. \tag{7.6}$$

Hence $A_E\xi = TE$ and $||A_E\xi||^2 = ||TE||^2 = \frac{s^2a^4}{4b^4}$. It follows that

$$K(P_{E\xi}) = \frac{s^2a^2}{4b^4},$$

where $K(P_{EF})$ denotes the sectional curvature of the plane generated by vectors E, F. If $E, F \in \mathcal{H}$ then

$$K(P_{EF}) = K_*(P_{E_*F_*}) - \frac{3g(E, TF)^2}{a^2||E \wedge F||^2},$$

where E_* denotes the projection of E on M i.e. $E_* = p(E)$. Thus

$$K(P_{EF}) = \frac{1}{b^2}K_0(P_{E_*F_*}) - \frac{3s^2a^2g(E, \tilde{J}F)^2}{4b^4||E \wedge F||^2}, \tag{7.7}$$

where K_0 stands for the sectional curvature of the metric h on N. Applying this we get for any $E \in \mathcal{H}$ the formula for the Ricci tensor ρ of (M, g):

$$\rho(E, E) = \frac{1}{b^2}\rho_0(bE_*, bE_*) - \frac{3s^2a^2}{4b^4} + \frac{s^2a^2}{4b^4},$$

where ρ_0 is a Ricci tensor of (M, h). Hence

$$\mu = \frac{\mu_0}{b^2} - \frac{s^2a^2}{2b^4},$$

where $\rho_0 = \mu_0 g_0$. Now we shall find a formula for $R(X, \xi)Y$ where $X, Y \in \mathcal{H}$. We have $R(X, \xi)Y = \nabla T(X, Y)$ and

$$\begin{aligned} \nabla T(X, Y) = \nabla_X(T(Y)) - T(\nabla_X Y) = \nabla^*_{X_*}(\tilde{T}Y^*) + \frac{1}{2}\mathcal{V}[X, TY] \\ -(\tilde{T}(\nabla^*_{X_*}Y_*))^* = \frac{1}{2}\mathcal{V}[X, TY] = -\frac{1}{2}sp^*\Omega_N(X, TY)\xi = -\frac{s^2a^2}{4b^2}h(X_*, Y_*)\xi \end{aligned} \tag{7.8}$$

Consequently $R(X, Y, Z, \xi) = 0$ for $X, Y, Z \in \mathcal{H}$,and

$$R(X, \xi, Y, \xi) = -\frac{s^2a^4}{4b^2}h(X_*, Y_*). \tag{7.9}$$

8. Riemannian submersion $p : (0, L) \times P \to (0, L)$. In this case the O'Neill tensor $A = 0$. We shall compute the O'Neill tensor T (see [ON]). Let us denote by Y^* the horizontal lift of the vector $Y \in TN$ with respect to the Riemannian submersion $p_N : P \to N$ i.e. $p_N(Y^*) = Y, g(Y^*, \xi) = 0$. Let $H = \frac{d}{dt}$ be the horizontal vector field for this submersion and $\mathcal{D}$ be the distribution spanned by the vector fields H, ξ. If $U, V \in \mathcal{V}$ and $g(U, V) = 0$ then $T(U, V) = 0$. Let $U \in \mathcal{V}, g(U, \xi) = 0$ and $U = U^*_*$ with $h(U_*, U_*) = 1$, then the following formula holds

$$T(U, U) = -rr'H. \tag{8.1}$$

In fact $2g(\nabla_U V, H) = -Hg(U, V) = -2rr'h(U_*, V_*)$ if $U = V$ or 0 if $g(U, V) = 0$. We also have

$$T(\xi, \xi) = -ff'H. \tag{8.2}$$

Now we shall prove that the almost complex structure defined by

$$JH = \frac{1}{f}\xi, JX = (J_*X_*)^* \text{ for } X = (X_*)^* \in \mathcal{E} = \mathcal{D}^{\perp}$$

where $X_* \in TN$, is a Kähler structure with respect to the metric g. We have for horizontal lifts $X, Y \in \mathfrak{X}(P) \subset \mathfrak{X}((0,L) \times P)$ of the fields $X_*, Y_* \in \mathfrak{X}(N)$ (with respect to the submersion described in the Section 7):

$$\begin{gathered}\nabla J(Y,X) = \nabla_Y(JX) - J(\nabla_Y X) = \nabla^*_{Y_*}(J_*(X_*))^* - \frac{1}{2}d\theta(Y,JX)\xi \\ +T(Y,JX) - J(\nabla^*_{Y_*}(X_*)^* - \frac{1}{2}d\theta(Y,X)\xi + T(Y,X)) = \\ -\frac{1}{2}sh(JY,JX)fJH - rr'h(Y,JX)H - \frac{1}{2}sh(JY,X)fH + h(X,Y)rr'JH = 0\end{gathered}$$

if and only if $f = \frac{2rr'}{s}$. Since the distribution $\mathcal{D}$ is totally geodesic and two-dimensional it is clear that $\nabla J(X,Y) = 0$ if $X, Y \in \Gamma(\mathcal{D})$. Now we shall show that

$$\nabla J(H,X) = \nabla J(X,H) = 0 \text{ for } X \in \mathcal{E}.$$

It is easy to show that $\nabla_X H = \nabla_H X = \frac{r'}{r}X$ and

$$\nabla_X(JH) = \nabla_X(\frac{1}{f}\xi) = \frac{1}{f}T(X) = \frac{sf}{2r^2}JX.$$

On the other hand

$$\nabla_X(JH) = \nabla J(X,H) + J(\nabla_X H) = \nabla J(X,H) + \frac{r'}{r}J(X).$$

Thus $\nabla J(X,H) = 0$ if $f = \frac{2rr'}{s}$. Similarly

$$\nabla_H(JX) = \nabla_{JX}H = \frac{r'}{r}JX = \nabla J(H,X) + J(\nabla_H X) = \nabla J(H,X) + \frac{r'}{r}JX$$

and $\nabla J(H,X) = 0$. Note that the Kähler form $\Omega = fdt \wedge \theta + r^2 p^*\Omega_N$ of almost Hermitian manifold $((0,L) \times P, g, J)$ is closed, which means that the structure J is almost Kähler. Thus $\nabla J(JX,Y) = -J\nabla J(X,Y)$ and consequently $\nabla_\xi J = 0$ which finishes the proof. Let $U, V, W \in \mathcal{V}$ and $g(U,\xi) = g(V,\xi) = g(W,\xi) = 0$. Then

$$R(U,V,\xi,W) = \hat{R}(U,V,\xi,W) - g(T(U,\xi),T(V,W)) + g(T(V,\xi),T(U,W)) = 0.$$

From O'Neill formulae it follows also that

$$R(JH,U,V,JH) = 0$$

if $g(U,V) = 0$ and

$$R(JH,U,U,JH) = \frac{s^2f^2}{4r^4} + \frac{f'r'}{fr}, \tag{8.3}$$

for a unit vector field U as above. Note also that the distribution $\mathcal{D}$ spanned by the vector fields ξ, H is totally geodesic. Consequently if $X, Y, Z \in \Gamma(\mathcal{D})$ and V is as above then

$$R(X, Y, Z, V) = 0. \tag{8.4}$$

Theorem 8.1. *The Kähler metric $g = dt^2 + (\frac{2rr'}{s})^2\theta^2 + r^2p^*h$ on the manifold $(0, L) \times P$ has quasi-constant holomorphic sectional curvature if and only if $dimN = 2$ or the Kähler manifold (N, h) has constant holomorphic sectional curvature and $dimN \geq 4$.*

Proof. Let $X = X^* + \alpha H + \beta JH$, where $X^* \in \mathcal{E} = \mathcal{D}^\perp$. Let us assume that $g(X, X) = 1$. Note that $|X_{\mathcal{D}}|^2 = \alpha^2 + \beta^2$ and $|X^*|^2 = 1 - |X_{\mathcal{D}}|^2$ and $JH = \frac{1}{f}\xi$ where $f = \frac{2rr'}{s}$. We also have

$$\begin{gathered}
R(X^* + \alpha H + \beta JH, JX^* + \alpha JH - \beta H, JX^* + \alpha JH - \beta H, X^* + \alpha H + \beta JH) = \\
a^4R(H, JH, JH, H) - a^2\beta^2R(H, JH, H, JH) + \beta^4R(JH, H, H, JH) \\
-\alpha^2\beta^2R(JH, H, JH, H) + \alpha^2R(H, JX^*, JH, X^*) + \alpha^2R(H, JX^*, JX^*, H) \\
+\alpha^2R(X^*, JH, JX^*, H) + \alpha^2R(X^*, JH, JH, X^*) + \beta^2R(JH, JX^*, -H, X^*) \\
\beta^2R(JH, JX^*, JX^*, JH) + \beta^2R(X^*, H, H, X^*) + \beta^2R(X^*, -H, JX^*, JH) \\
+2\alpha^2R(H, JH, JX^*, X^*) - 2\beta^2R(JH, H, JX^*, X^*) \\
+R(X^*, JX^*, JX^*, X^*) = |X_{\mathcal{D}}|^4R(H, JH, JH, H) + 8|X_{\mathcal{D}}|^2R(H, JX^*, JX^*, H) \\
+R(X^*, JX^*, JX^*, X^*) = |X_{\mathcal{D}}|^4R(H, JH, JH, H)+ \\
8|X_{\mathcal{D}}|^2(1 - |X_{\mathcal{D}}|^2)R(JH, \tilde{X}^*, \tilde{X}^*, JH) + (1 - |X_{\mathcal{D}}|^2)^2R(\tilde{X}^*, J\tilde{X}^*, J\tilde{X}^*, \tilde{X}^*)
\end{gathered}$$

where $\tilde{X}^* = \frac{1}{|X^*|}X^*$ if $X^* \neq 0$ or $\tilde{X}^* = 0$ if $X^* = 0$.

From (8.5) it follows that $R(X^* + \alpha H + \beta JH, JX^* + \alpha JH - \beta H, JX^* + \alpha JH - \beta H, X^* + \alpha H + \beta JH)$ depends only on $|X_{\mathcal{D}}|$ and the point $x \in (0, L) \times P$ if and only if $R(\tilde{X}^*, J\tilde{X}^*, J\tilde{X}^*, \tilde{X}^*)$ does not depend on the unit vector $\tilde{X}^*$. It follows that $dimN = 2$ or $dimN \geq 4$ and at every point $x \in N$ the holomorphic sectional curvature of (N, h) is constant. In fact from O'Neill formulae ([ON]) it follows that

$$\begin{gathered}
R(\tilde{X}^*, J\tilde{X}^*, J\tilde{X}^*, \tilde{X}^*) = R_*(\tilde{X}^*, J\tilde{X}^*, J\tilde{X}^*, \tilde{X}^*) - g(T(\tilde{X}^*, \tilde{X}^*), T(J\tilde{X}^*, J\tilde{X}^*)) \\
+2g(A(\tilde{X}^*, J\tilde{X}^*), A(J\tilde{X}^*, \tilde{X}^*)) - g(A(J\tilde{X}^*, \tilde{X}^*), A(J\tilde{X}^*, \tilde{X}^*)) \\
= R_*(\tilde{X}^*, J\tilde{X}^*, J\tilde{X}^*, \tilde{X}^*) - 4\frac{(r')^2}{r^2} = \frac{c_0}{r^2} - 4\frac{(r')^2}{r^2},
\end{gathered}$$

where we used the formula $A(E, F) = \frac{s}{2r^2}g(E, \tilde{J}F)\xi$. Note that if (N, h, J) has constant holomorphic sectional curvature c_0 then the coefficient a in the formula (2.1) for the tensor R equals $a = \frac{c_0}{r^2} - 4\frac{(r')^2}{r^2}$ and is non-constant. Note that $div_{\mathcal{E}}H = 2(n-1)\frac{r'}{r}$ and $div_{\mathcal{E}}\xi = 0$. In particular $\kappa = 2(n-1)\frac{r'}{r} \neq 0$ on an open and dense subset. ♢

Remark. Note that in the case $s = 0$ we also get a Kähler manifold with QCH metric and a special Killing Ricci potential. However in this case $\kappa = 0$ on the whole of $(0,L)\times P$. Let g_0 be any smooth, Riemannian metric on $\mathbb{CP}^1$. If h is a metric of constant holomorphic sectional curvature on $\mathbb{CP}^n$, then the product $(\mathbb{CP}^1\times\mathbb{CP}^{n-1}, g_0\times h)$ is a compact, simply connected manifold QCH Kähler manifold (with respect to the distribution whose leaves are submanifols $\mathbb{CP}^1\times\{x\}$, for $x\in\mathbb{CP}^{n-1}$), having $\kappa=0$ and, in general, not admitting a special Kähler-Ricci potential.

9. QCH Kähler manifolds with real analytic Riemannian metric. We start with :

Lemma 9.1. *Let us assume that (M,g,J) is a (connected, complete) QCH Kähler manifold with a real analytic metric g and $\dim M\geq 6$. Then $int\ B=\emptyset$ or $b=0$ and (M,g,J) has a constant holomorphic sectional curvature.*

Proof. We can find local coordinates $(x_1,x_2,...,x_{2n})$ such that $g_{\alpha\beta}$ are real analytic, where $g_{\alpha\beta}=g(\partial_\alpha,\partial_\beta)$ and $\partial_\alpha=\frac{\partial}{\partial x_\alpha}$. Analogously $\rho_{\alpha\beta}=\rho(\partial_\alpha,\partial_\beta)$ are real analytic functions. If $g(SX,Y)=\rho(X,Y)$, where ρ is the Ricci tensor of (M,g,J), then $S_\alpha^\beta=g^{\gamma\beta}\rho_{\alpha\gamma}$ where $S\partial_\alpha=S_\alpha^\beta\partial_\beta$. It follows that the functions S_α^β are analytic. Consequently the polynomial $\phi(t)=det(S-tI)$ has real analytic coefficients. Thus if $b=0$ in open set $V\subset U$ then also $c=0$ on V and $S=\lambda I$ on V for a constant $\lambda\in\mathbb{R}$. This means that $\phi(t)=(\lambda-t)^{2n}$ in V and consequently on the whole of M. Thus $S=\lambda I$ on the whole of M. It follows (see (2.13)) that $\frac{n+1}{2}a+\frac{b}{4}$ is constant on the whole of M and $\frac{n+2}{4}b+c=0$ on the whole of M. Consequently $b+4c=-(n+1)b$. From (2.9) we get in $U-B$:

$$\xi(\ln|b|)=-\frac{n+1}{n-1}\kappa \tag{9.1}$$

Let $t\to c(t)$ be a unit geodesic joining a point $c(0)=x\in U-B$ with a point $c(l)=y\in B$. Note that $|\dot c(\ln|b|)|\leq|\xi(\ln|b|)|=\frac{n+1}{n-1}\kappa$. Thus we obtain $\ln|b|\circ c(t)-\ln|b|\circ c(0)\geq-\frac{n+1}{n-1}\int_0^t\kappa\circ c(s)ds\geq-\frac{n+1}{n-1}\kappa_0 l$ where $0\leq\kappa\circ c(s)\leq\kappa_0=\sup_{c([0,l])}\kappa$. It is clear now that $U-B=\emptyset$ and $b=0$ in U. It follows that in the case of analytic metric the assumption $int\ B\neq\emptyset$ implies $b=0,c=0$ on the whole of U and hence on M which means that (M,g,J) has constant holomorphic sectional curvature.$\diamondsuit$

Now we shall prove

Theorem 9.2. *Let (M,g,J) be a compact, simply connected QCH Kähler manifold of dimension $2n\geq 6$. If $\kappa\neq 0$ and $int\ B=\emptyset$ then M is a $\mathbb{CP}^1$-bundle over $\mathbb{CP}^{n-1}$ with a metric homothetic to the metric (6.1) where $s\neq 0, f=\frac{2rr'}{s}, s=\frac{2k}{n}, k\in\mathbb{N}$, r satisfies the boundary conditions (6.4) and h is a Kähler metric of constant holomorphic sectional curvature on $\mathbb{CP}^{n-1}$.*

Proof. From Theorem 5.3 it follows that (M,g,J) is a compact Kähler manifold admitting a special Killing-Ricci potential. From the classification of such manifolds given by Derdziński and Maschler (see [D-M-1]) it follows that M is a holomorphic $\mathbb{CP}^1$-bundle over a compact, simply connected Einstein-Kähler manifold (N,h), i.e. r is given by formulae (6.1) or (6.2) with f,r satisfying the initial conditions. Since $\kappa\neq 0$ it follows that $s\neq 0$ and the metric is homethetic to the metric (6.1) with $f=\frac{2rr'}{s}$ where r is a smooth function, positive on $(0,L)$, even at $0,L$ and with $r'>0$ satisfying the boundary conditions (6.4). As (M,g,J) is a QCH Kähler manifold it follows that (N,h) is compact, simply connected Kähler manifold of constant holomorphic curvature, which means that $N = \mathbb{CP}^{n-1}$ with a standard Fubini-Study metric.$\diamondsuit$

Theorem 9.3. *Let (M, g, J) be a compact, simply connected QCH Kähler manifold of dimension $2n \geq 6$ with an analytic Riemannian metric g. If $\kappa \neq 0$ then M is a $\mathbb{CP}^1$-bundle over $\mathbb{CP}^{n-1}$ with a metric homothetic to the metric (6.1) where $s \neq 0, f = \frac{2rr'}{s}, s = \frac{2k}{n}, k \in \mathbb{N}$ with $r \in C^{\omega}(\mathbb{R})$, r satisfies the boundary conditions (6.4) and h is a Kähler metric of constant holomorphic sectional curvature on $\mathbb{CP}^{n-1}$ with scalar curvature $4(n-1)$ or (M, g, J) is a projective complex space $\mathbb{CP}^n$ with a metric homothetic to the standard Fubini-Study metric.*

Proof. If the metric g is analytic then $int\ B = \emptyset$ or $\{x : b(x) = 0, c(x) = 0\} = M$. In the first case we apply Theorem 9.2 and in the second case (M, g, J) has constant holomorphic sectional curvature at every point $x \in M$ which means that (M, g, J) has constant holomorphic sectional curvature (see [K-N], Chapter 9, Theorem 7.5, p.158 of Russian translation). Since M is compact and simply connected it follows (see [K-N], Chapter 9, Theorems 7.8, 7.9 pp.160-161) that $M = \mathbb{CP}^n$ with a metric homothetic to the standard Fubini-Study metric. $\diamondsuit$

Remark. Note, that the tangent bundle $T\mathbb{CP}^n$ may not admit any complex line subbundle. This is the case if $n = 2$ ($T\mathbb{CP}^2$ does not contain any two-dimensional, oriented real subbundle, see [H-H]). Thus $\mathbb{CP}^2$ does not admit any J-invariant two-dimensional global distribution $\mathcal{D}$ and cannot be a QCH Kähler manifold.

Now we show that there are uncountably many analytic functions r satisfying the boundary conditions (6.4). Note first that the function $(r')^2$ is a function of r alone, i.e. $(r')^2 = P(r)$ for a certain function P which is smooth if r is smooth. We shall give a family of analytic functions P which are polynomials of degree 3, parameterized by the real numbers x, y where $im\ r = [x, y]$ which give rise to functions r satisfying the boundary conditions (6.4) and thus giving examples of QCH Kähler manifolds. In that way we show one of the method of constructing such functions. Let $0 < x < y$ and let us consider a polynomial

$$(9.2)\qquad P(t) = -\frac{s(x+y)}{y-x} + \frac{s(x^2+3sxy+y^2)}{xy(y-x)}t - \frac{2s(x+y)}{xy(y-x)}t^2 + \frac{s}{xy(y-x)}t^3.$$

Then $P(x) = P(y) = 0$ and $P(t) > 0$ for $t \in (x, y)$ and

$$(9.3)\qquad P'(x)x = s, P'(y)y = -s.$$

Let us consider the ordinary differential equation

$$(9.4)\qquad r'' = \frac{1}{2}P'(r)$$

with the initial conditions $r(0) = x, r'(0) = 0$. Then any solution r of (9.4) is an analytic function $r \in C^{\omega}(\mathbb{R})$. Let $L > 0$ be the first point such that $r(L) = y$. Then for $t \in (0, L)$ we have

$$(9.5)\qquad r'(t) = \sqrt{P(r)}.$$

Note that

$$L=\int_x^y \frac{dt}{\sqrt{P(t)}}. \tag{9.6}$$

The function r is even at both points $0, L$. In fact we have $r'(0)=0, r'(L)=0$ and r satisfies (9.4). It follows that $r^{(2k-1)}(0)=r^{(2k-1)}(L)=0$ for $k\in\mathbb{N}$. Since $r\in C^{\omega}(\mathbb{R})$ it follows that r is even at $0, L$. In view of (9.5) $r'(t)>0$ for every $t\in(0,L)$. We also have

$$2r(0)r''(0)=2x\frac{1}{2}P'(x)=xP'(x)=s$$
$$2r(L)r''(L)=2y\frac{1}{2}P'(y)=yP'(y)=-s.$$

It follows that the metric $g=dt^2+(\frac{2rr'}{s})^2\theta^2+r^2p^*h$ on $(0,L)\times P$ where L is given by (9.6) and r is the solution of (9.4) extends to an analytic metric on the $\mathbb{CP}^1$-bundle $M=P\times_{S^1}\mathbb{CP}^1$. Note that the special Killing-Ricci potential is $\tau=\frac{r^2}{s}$ and the holomorphic Killing vector field corresponding to it is ξ with $g(\xi,\xi)=f^2=(\frac{2rr'}{s})^2$. We also have $\kappa=2(n-1)\frac{r'}{r}$.

Theorem 9.4. *Let (M,g,J) be a complete QCH Kähler manifold of dimension $2n\geq 6$ and an analytic Riemannian metric g. If $\kappa\neq 0$ then int $B=\emptyset$, the set $U=\{x\in M:\kappa(x)\neq 0\}$ is open and dense in M and every point $x\in U$ has a neighborhood $V\subset U$ biholomorphic to the manifold $(\alpha,\beta)\times P$ with the metric (6.1) where $s\neq 0, f=\frac{2rr'}{s}, s\in\mathbb{R}$ with $r\in C^{\omega}(\mathbb{R})$, P is a circle bundle over a Kähler manifold (N,h) and h is a Kähler metric of constant holomorphic sectional curvature or $b=0$ and M has a metric of constant holomorphic sectional curvature.*

Proof. The proof follows from Section 3, Lemma 9.1, Theorem 5.1 and from [D-M-2] (see Theorem 18.1).$\diamondsuit$

Acknowledgments. The author would like to thank the referee for his valuable remarks which improved the paper.

References.

[Bes] A. L. Besse *Einstein manifolds*, Ergebnisse, ser.3, vol. 10, Springer-Verlag, Berlin-Heidelberg-New York, 1987.

[Ber] L. Bérard Bergery,*Sur de nouvelles variétés riemanniennes d'Einstein*, Publ. de l'Institute E. Cartan (Nancy) **4**,(1982), 1-60.

[D-M-1] A. Derdziński, G. Maschler *Special Kähler-Ricci potentials on compact Kähler manifolds*, J. reine angew. Math. 593 (2006), 73-116.

[D-M-2] A. Derdziński, G. Maschler *Local classification of conformally-Einstein Kähler metrics in higher dimensions*, Proc. London Math. Soc. (3) 87 (2003), no. 3, 779-819.

[G-M-1] G.Ganchev, V. Mihova *Kähler manifolds of quasi-constant holomorphic sectional curvatures*, Cent. Eur. J. Math. 6(1),(2008), 43-75.

[G-M-2] G.Ganchev, V. Mihova *Warped product Kähler manifolds and Bochner-Kähler metrics*, J. Geom. Phys. 58(2008), 803-824.

[H-H] F. Hirzebruch, H. Hopf *Felder von Flächenelementen in 4-dimensionalen Mannigfaltigkeiten.*, Math. Ann. 136, (1958), 156-172.

[K] S. Kobayashi *Principal fibre bundles with the 1-dimensional toroidal group* Tôhoku Math.J. **8**,(1956) 29-45.

[K-N] S. Kobayashi and K. Nomizu *Foundations of Differential Geometry*, vol.2, Interscience, New York 1963

[ON] B. O'Neill, *The fundamental equations of a submersion*, Mich. Math. J. **13**,(1966), 459-469.

[S] P. Sentenac ,*Construction d'une métrique d'Einstein sur la somme de deux projectifs complexes de dimension 2*, Géométrie riemannienne en dimension 4 (Séminaire Arthur Besse 1978-1979) Cedic-Fernand Nathan, Paris (1981), pp. 292-307.

Institute of Mathematics
Cracow University of Technology
Warszawska 24
31-155 Kraków, POLAND.
E-mail address: wjelon@pk.edu.pl